



KCNA1 gene

potassium voltage-gated channel subfamily A member 1

Normal Function

The *KCNA1* gene belongs to a large family of genes that provide instructions for making potassium channels. These channels, which transport positively charged atoms (ions) of potassium into and out of cells, play a key role in a cell's ability to generate and transmit electrical signals.

The *KCNA1* gene provides instructions for making one part (the alpha subunit) of a potassium channel called Kv1.1. These channels are found in the brain, where they transport potassium ions into nerve cells (neurons). The flow of certain ions, including potassium, into and out of neurons regulates communication between these cells.

Health Conditions Related to Genetic Changes

episodic ataxia

At least 20 mutations in the *KCNA1* gene have been identified in people with episodic ataxia type 1 (EA1). People with this form of the condition have brief, recurrent episodes of poor coordination and balance (ataxia). Between episodes, many affected individuals experience myokymia, a muscle abnormality that can cause involuntary muscle cramping, stiffness, and continuous, fine muscle twitching that appears as rippling under the skin.

Most of the *KCNA1* mutations responsible for episodic ataxia change single protein building blocks (amino acids) in the alpha subunit of the Kv1.1 channel. Some of these changes prevent the assembly of functional channels, while other mutations alter the channel's structure. When Kv1.1 channels are missing or abnormal, the flow of potassium ions into neurons is reduced. This decrease in potassium ions overexcites certain neurons in the brain, which disrupts normal communication between these cells. Although changes in signaling between neurons underlie the episodes of uncoordinated movement seen in people with episodic ataxia, it is unclear how altered potassium ion transport causes the specific features of the condition.

other disorders


Mutations in the *KCNA1* gene have been found to cause a range of signs and symptoms affecting the nervous system. In at least one family, isolated myokymia (continuous muscle twitching and spasms without episodes of ataxia) has been attributed to *KCNA1* mutations. Changes in this gene have also been identified in

a small number of people with epilepsy. Like the *KCNA1* mutations that underlie episodic ataxia, the mutations that cause isolated myokymia and epilepsy reduce the flow of potassium ions through Kv1.1 channels, disrupting normal communication between neurons in the brain. Researchers are working to determine why mutations in this single gene can cause several different disorders of the nervous system.

Chromosomal Location

Cytogenetic Location: 12p13.32, which is the short (p) arm of chromosome 12 at position 13.32

Molecular Location: base pairs 4,909,907 to 4,918,256 on chromosome 12 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- AEMK
- EA1
- HBK1
- HUK1
- KCNA1_HUMAN
- KV1.1
- MBK1
- MGC126782
- MGC138385
- MK1
- potassium channel, voltage gated shaker related subfamily A, member 1
- potassium voltage-gated channel, shaker-related subfamily, member 1 (episodic ataxia with myokymia)

- RBK1
- voltage-gated potassium channel subunit Kv1.1

Additional Information & Resources

Educational Resources

- Biochemistry (fifth edition, 2002): Specific Channels Can Rapidly Transport Ions Across Membranes
<https://www.ncbi.nlm.nih.gov/books/NBK22509/>
- Neuromuscular Disease Center, Washington University: Potassium channels
<http://neuromuscular.wustl.edu/mother/chan.html#k>

GeneReviews

- Episodic Ataxia Type 1
<https://www.ncbi.nlm.nih.gov/books/NBK25442>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28KCNA1%5BTIAB%5D%29+OR+%28%28HBK1%5BTIAB%5D%29+OR+%28KV1.1%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- POTASSIUM CHANNEL, VOLTAGE-GATED, SHAKER-RELATED SUBFAMILY, MEMBER 1
<http://omim.org/entry/176260>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_KCNA1.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=KCNA1%5Bgene%5D>
- HGNC Gene Family: Potassium voltage-gated channels
<http://www.genenames.org/cgi-bin/genefamilies/set/274>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=6218

- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/3736>
- UniProt
<http://www.uniprot.org/uniprot/Q09470>

Sources for This Summary

- Browne DL, Gancher ST, Nutt JG, Brunt ER, Smith EA, Kramer P, Litt M. Episodic ataxia/myokymia syndrome is associated with point mutations in the human potassium channel gene, KCNA1. *Nat Genet.* 1994 Oct;8(2):136-40.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/7842011>
- Chen H, von Hehn C, Kaczmarek LK, Ment LR, Poher BR, Hisama FM. Functional analysis of a novel potassium channel (KCNA1) mutation in hereditary myokymia. *Neurogenetics.* 2007 Apr;8(2):131-5. Epub 2006 Nov 29.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17136396>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1820748/>
- Eunson LH, Rea R, Zuberi SM, Youroukos S, Panayiotopoulos CP, Liguori R, Avoni P, McWilliam RC, Stephenson JB, Hanna MG, Kullmann DM, Spauschus A. Clinical, genetic, and expression studies of mutations in the potassium channel gene KCNA1 reveal new phenotypic variability. *Ann Neurol.* 2000 Oct;48(4):647-56.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11026449>
- Jen JC, Graves TD, Hess EJ, Hanna MG, Griggs RC, Baloh RW; CINCH investigators. Primary episodic ataxias: diagnosis, pathogenesis and treatment. *Brain.* 2007 Oct;130(Pt 10):2484-93. Epub 2007 Jun 15. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17575281>
- Maylie B, Bissonnette E, Virk M, Adelman JP, Maylie JG. Episodic ataxia type 1 mutations in the human Kv1.1 potassium channel alter hKvbeta 1-induced N-type inactivation. *J Neurosci.* 2002 Jun 15;22(12):4786-93.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12077175>
- Rajakulendran S, Schorge S, Kullmann DM, Hanna MG. Episodic ataxia type 1: a neuronal potassium channelopathy. *Neurotherapeutics.* 2007 Apr;4(2):258-66. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17395136>
- Rea R, Spauschus A, Eunson LH, Hanna MG, Kullmann DM. Variable K(+) channel subunit dysfunction in inherited mutations of KCNA1. *J Physiol.* 2002 Jan 1;538(Pt 1):5-23.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11773313>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2290030/>
- Zerr P, Adelman JP, Maylie J. Episodic ataxia mutations in Kv1.1 alter potassium channel function by dominant negative effects or haploinsufficiency. *J Neurosci.* 1998 Apr 15;18(8):2842-8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/9526001>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/KCNA1>

Reviewed: August 2008
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services